

Introduction to Botany. Lecture 25

Alexey Shipunov

Minot State University

November 6, 2015



1 Questions and answers

2 Root

- Root morphology
- Anatomy and development of roots
- Origins of root tissues
- Water transport in roots
- Diversity of roots



1 Questions and answers

2 Root

- Root morphology
- Anatomy and development of roots
- Origins of root tissues
- Water transport in roots
- Diversity of roots



Previous final question: the answer

What is angle of divergence?



Previous final question: the answer

What is angle of divergence?

- Angle between orthostychy: radial planes in which leaves are located around stem.




Root

Root morphology



Root system and shoot system



© 2006 Brooks/Cole - Thomson



Definition and functions

- Axial vegetative organ with a function of soil nutrition
- Other functions:
 - ① Anchor
 - ② Synthesis
 - ③ Storage
 - ④ Communication
- Features:
 - ① No leaves
 - ② Geotropic growth
 - ③ Locates in soil or water



Types of roots

- Primary root: originates from root of seedling
- Secondary (lateral) roots: originate from primary roots
- Adventitious roots: originate from stems



Primary root



Adventitious roots




Root systems

- Tap root system: with well developed primary root (most seed plants)
- Fibrous root system: without clearly visible primary root (monocots, ferns)



Fibrous and tap root systems



© 2006 Brooks/Cole - Thomson




Root

Anatomy and development of roots



Root zones




- Root cap
- Root meristem
- Elongation zone
- Absorption zone
- Maturation zone

Structure of root tip

- Initial cells (quiescent center)
- RAM
- Root tip growing both forward (root cap) and backward (other root tissues), initial cells determine the direction of growth
- If root tip touch barrier, it starts to make rotating movements



Root tip



© 2006 Brooks/Cole - Thomson



Periphery of root

- Rhizoderm (root epidermis): fast-degrading cells
- Cortex, which includes also:
 - Endoderm: 1-cell layer with specialized cell walls, located on the border with vascular cylinder
 - And (sometimes) exoderm: similar to endoderm but located just under rhizoderm
- In some plants (i.e., orchids), cortex modified into velamen




Root center: vascular cylinder

- Pericycle
- Vascular tissues located in the center
- No central hollow, central parenchyma presents in monocot roots



Anatomy of root



© 2006 Brooks/Cole - Thomson




Pericycle

- Long-lived parenchyma cells served as half-meristem
- Initiates development of lateral roots
- Contributes to vascular cambium
- Contributes to cork cambium



Development of lateral roots




Vascular bundle


- Only one!
- Has radial (star-like) symmetry
- Xylem arranged in rays, multiple in monocots, 2-4 in other plants



Radial structure of root vascular bundle in buttercup (*Ranunculus* sp.)



Root of monocot (*Zea mays*)




Root

Origins of root tissues



Development of tissues



© 2006 Brooks/Cole - Thomson



Root

Water transport in roots



Rhizoderm and osmosis

- The existence of root hairs dramatically increases the surface of absorption
- Every root hair cell increase the internal concentration of large molecules, typically organic acids
- Process of concentration requires ATP
- As a result, osmosis water flow starts from soil to root cells




Endoderm and root pressure


- From rhizoderm to endoderm, transport of water is both symplastic and apoplastic
- In the endoderm cells, Caspari stripes stop apoplastic transport and therefore forced symplastic transport
- This is a high-energetic process requires ATP
- As a result, water will be pushed up from root: this is the root pressure




Apoplastic and symplastic transport in the root



Casparian strips




© 2006 Brooks/Cole - Thomson



© 2006 Brooks/Cole - Thomson




How Casparian strips are working



© 2006 Brooks/Cole - Thomson



Origin of root pressure



Root

Diversity of roots




Modifications of roots

- Adventive buds with root origin (many plants)
- Mycorrhizae: endotrophic (grasses, orchids) and ectotrophic (trees)
- Haustoria (parasites like *Cuscuta*—dodder plant)
- Root nodules (legumes, Fabaceae family)
- Contractile roots (*Hyacinthus* spp.—hyacinth, *Taraxacum* spp.—dandelion)
- Storage roots (*Daucus carota*—carrot, *Armoracia officinalis*—horseradish)
- Supportive roots (many tropical plants)
- Defensive, spiny roots (ivy)
- Photosynthetic roots (some orchids)




Endotrophic mycorrhizae in *Corallorhiza* orchid




© 2006 Brooks/Cole - Thomson



Ectotrophic mycorrhizae of trees



Haustoria of *Cuscuta* (dodder)



Nodulated roots of soybean (*Glycine max*)



Contractile roots of *Hyacinthus orientalis*



© 2006 Brooks/Cole - Thomson



Supportive roots of mangrove plants



Supportive roots of *Pandanus* sp.



Defensive spiny roots of ivy (*Hedera* sp.)



Photosynthetic aerial roots of orchids



Table of modifications

Function	Stem	Leaf	Root
Expansion	...	Plantlets	Adventive buds
Storage	...	Succulent leaves	Storage roots
Photosynthesis	...	DEFAULT	Some aerial roots
Defense	...	Spines, scale	Root spines
Support	DEFAULT	Leaf ten-drills	Haustoria, aerial and contractile roots
Interactions	...	Traps, "sticky tapes", urns	Mycorrhizae, nodulated roots



For Further Reading



A. Shipunov.

Introduction to Botany [Electronic resource].

2015.

Mode of access:

http://ashipunov.info/shipunov/school/biol_154

