

Biometry. Lecture 16

Alexey Shipunov

Minot State University

April 2, 2014



- 1 Questions and answers
- 2 Two-dimensional statistics
 - Regression



- 1 Questions and answers
- 2 Two-dimensional statistics
 - Regression



Starting...

```
> setwd("<working folder>")  
or  
"Change dir"  
in menu!
```



Two-dimensional statistics

Regression



The idea of regression

- The basic regression analysis will apply the linear model for data
- It will study not only if variables are associates and not only the strength of association but also the form of association (law of association)



Regression formula

- The simplest is a linear regression,

$$m = b_0 + b_1 \times x,$$

where m is a **predicted value**, x is an **independent variable** and b_0 and b_1 are coefficients (so-called **intersect** and **slope**).

- In other terms, linear regression is
response = intersect + slope * influence
- In R model formula language, it is simply
response ~ influence



Analysis of regression model

- If y is a real response, then error of model

$$E = y - m$$

- If σ^2 are dispersions of m and y , then

$$R^2 = 1 - \sigma_m^2 / \sigma_y^2,$$

- In a background, R^2 is similar to coefficient of determination



Test of regression

- To test if regression model is correct, the Fisher test is normally applied
- Null hypothesis for Fisher test is that a model is not reliable



Regression example: women data

```
> lm.women <- lm(weight ~ height, data = women)
> plot(weight ~ height, data = women, main="",
+ xlab="Height (feet)", ylab="Weight (pounds)")
> grid()
> abline(lm.women, col="red")
```



Analysis of regression

```
> summary(lm.women)
```



Analysis of analysis

- Resulted model: $\text{weight} = -87.51667 + 3.45 * \text{height}$
- Maximum deviations from model are -1.7333 and 3.1167 pounds
- Almost half of residuals are between first and third quartiles
- All coefficients are significant
- Adjusted R-squared is close to 1 (very high!)
- The overall p-value is much less than 0.05 therefore the model is reliable
- There are 1 and 13 degrees of freedom (for columns and for rows)



Thuesen data example

- 24 rows and 2 columns data for observations of ventricular velocity with different levels of blood glucose
- Data was taken from patients with diabetes type I.



Running the example and explaining results

```
> install.packages("ISwR")
> library(ISwR)
> str(thuesen); head(thuesen)
> thuesen <- na.omit(thuesen)
> thuesen.lm <- lm(short.velocity ~ blood.glucose,
+ data=thuesen)
> thuesen.lm
> summary(thuesen.lm)
```



Scatterplot with regression line

```
> plot(short.velocity ~ blood.glucose, data=thuesen)
> abline(thuesen.lm)
```



Visualizing residuals

```
> with(thuesen, segments(blood.glucose,  
+ fitted(thuesen.lm), blood.glucose, short.velocity))
```



Confidence intervals for regression

```
> pred.frame <- data.frame(blood.glucose=4:20)
> pc <- predict(thuesen.lm, int="c", newdata=pred.frame)
> plot(short.velocity ~ blood.glucose, data=thuesen)
> pred.gluc <- pred.frame$blood.glucose
> matlines(pred.gluc, pc, lty=c(1,2,2), col="black")
```



Diagnostic plots for regression

- “Residuals vs. Fitted”: checks outliers, the best is flat line
- “Normal Q-Q”: checks residuals for normal distribution, if they are not normal then our regression is not linear
- “Scale-Location”: checks the trend in dispersion
- “Residuals vs. Leverage & Cook’s distance”: checks the most influential observations



Regression diagnostics

```
> plot(thuesen.lm)
> plot(lm(height ~ weight, data=women))
```



Finishing...

```
> savehistory("20140402.r")
```



Final question (3 points)



Final question (3 points)

In the embedded data `USArrests`, there are numbers of murders and rapes per 100,000 for every state. Are murders and rapes correlated? Is that correlation significant?



Summary: most important commands

- `lm()` —estimate the linear regression
- `predict()` —predict values with model



For Further Reading



A. Shipunov.

Biometry [Electronic resource].

2012—onwards.

Mode of access:

http://ashipunov.info/shipunov/school/biol_240



A. Shipunov, and others.

Visual statistics. Use R!

DMK Press, 2012. [Under translation from Russian.]

